



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\frac{f(k)}{10^{k-1}} (1 + r + r^2 + \dots).$$

This series converges, for r is < 1 ; and hence our given series converges.

The case of "improper" combinations may be rapidly dealt with by the following device. Write the digit a , k times at the end of the given combination, where a is any digit other than the first digit of the combination. This certainly gives us a "proper" combination (of $2k$ digits), so that, as already proved, the series formed from the harmonic series by omitting terms whose denominators contain this combination converges. But this series evidently contains all terms of the given series; this latter, then, converges too.

In like manner we may prove that *the series obtained by omitting terms containing all of a given set of combinations (some of which may be repeated) converges*. Here we may compare the series with that obtained by omitting terms containing the single combination formed from the given combinations by juxtaposition.

This supplies another proof of the proposition in my paper on "A Curious Convergent Series" in this MONTHLY for May, 1916, that proposition being the special case of the above in which each of the combinations consists of a single digit.

The proposition stated by Professor Moulton in his solution of Algebra 453 in this MONTHLY for Oct., 1916, refers, I suppose—the proof is not given in full—not to combinations of the kind here considered, but to selections, and without any reference to the juxtaposition of the digits of the selection in the omitted terms.

239 (Number Theory) [March, 1916; March, 1919]. Proposed by HAROLD T. DAVIS, Colorado Springs, Colorado.

Give a general method for determining the solution in integers of the equation

$$x^r - 10xy - (n + 1) + y = 0,$$

where r and n are positive integers.

II. SOLUTION BY FRANK IRWIN, University of California.

If $x = 0$, $y = n + 1$; noting this solution, we shall suppose in what follows $x \neq 0$.

The degree in x of the given equation may be reduced by the following device. $y - (n + 1)$ must be divisible by x , say $y - (n + 1) = y_1x$. Then our equation reduces to

$$x^{r-1} - 10xy_1 - 10(n + 1) + y_1 = 0.$$

Putting $y_1 - 10(n + 1) = y_2x$, $y_2 - 10^2(n + 1) = y_3x$, etc., we finally reach the equation

$$1 - 10xy_r - 10^r(n + 1) + y_r = 0, \quad (1)$$

where y and y_r are connected, it will be found, by the relation

$$y = (n + 1)[1 + 10x + 10^2x^2 + \dots + 10^{r-1}x^{r-1}] + y_rx^r,$$

or

$$y = (n + 1) \frac{10^rx^r - 1}{10x - 1} + y_rx^r. \quad (2)$$

Rewriting (1) as

$$(10x - 1)y_r = 1 - 10^r(n + 1), \quad (3)$$

we see that x and y_r must have opposite signs.

(i) Let y_r be negative, $y_r = -y'_r$. Then (3) becomes $(10x - 1)y'_r = 10^r(n + 1) - 1$. Here the number on the right ends with the digit 9, as does also the first factor on the left. Hence we see that y'_r must end with 1. We get, then, a solution by taking for y'_r any factor, ending with 1, of the right side, and determining x and y from (3) and (2).

(ii) Let x be negative, $x = -x'$. Then (3) becomes

$$(10x' + 1)y_r = 10^r(n + 1) - 1.$$

We take y_r equal to any factor of the right side that ends with 9.

This gives us all solutions.

Suppose that $10^r(n + 1) - 1$ can be factored as $a \cdot b$, where a ends with 1, b with 9. Then

we may take either $y_r = -a$, whence $x = (b+1)/10$; or $y_r = b$, $x = (-a+1)/10$. Thus solutions go in pairs, the x of either solution being obtained by adding 1 to the y_r of the other and dividing by 10. There are, then, always an even number of solutions, and always at least two, viz., those where $y_r = -1$, $x = 10^{r-1}(n+1)$; $y_r = 10^r(n+1) - 1$, $x = 0$.

Finally it may be noted that the more general equation

$$ax^r - bxy + y - c = 0$$

may be solved by similar methods. Here we should have to pick out such divisors of $b^r c - a$ as are $\equiv \pm 1 \pmod{b}$. It may be interesting to apply the method to an example. Let us solve

$$x^4 - 10xy - 22 + y = 0.$$

We have to solve

$$(10x - 1)y_4 = -219999 = -3 \times 13 \times 5641$$

(5641 prime). The values of y_4 with the corresponding solutions of our equation are:

$$y_4 = -5641, \quad x = 4, \quad y = 6; \quad y_4 = -1, \quad x = 22000 \text{ (} y \text{ a very large number)};$$

$$y_4 = 39, \quad x = -564, \quad y = -17937434; \quad y_4 = 219999, \quad x = 0, \quad y = 22.$$

247 (Number Theory) [June, 1916]. Proposed by NORMAN ANNING, Chilliwack, B. C.


To dissect the triangle whose sides are 52, 56, 60 into three Heronian triangles by lines drawn from the vertices to a point within.

The word Heronian is used in the sense of the German Heronische (Wertheim, *Anfangsgründe d. Zahlenlehre*, p. 140) to describe a triangle whose sides and area are integral.

SOLUTION BY FRANK IRWIN, University of California.

The orthocenter, O , may be taken as the required point. Let ABC be the triangle with $a = 60$, $b = 52$, $c = 56$; and let the feet of the perpendiculars from A, B, C on the opposite sides be D, E, F , respectively. Then the various lines in the figure, calculated as indicated below, are: $BD = 168/5$, $DC = 132/5$, $CE = 396/13$, $EA = 280/13$, $AF = 20$, $FB = 36$; $AO = 25$, $BO = 39$, $CO = 33$. Finally, area $BOC = 594$, area $COA = 330$, and area $AOB = 420$; so that the sides and areas of these three triangles are integral, as asserted.

The explanation of these facts depends on the following proposition: If the sides and area of the triangle ABC are rational, the same is true of the triangles BOC, COA, AOB . (Then by multiplying the dimensions of the figure by a suitable integer everything can be made integral.) For the three altitudes are rational, as also the radius r of the inscribed circle (since $rs = \text{area}$). Thus $\tan A/2$ is rational, and so, then, are $\cos^2 A/2$ and $\cos A$. Therefore, $AF = b \cos A$ is rational, and similarly, FB, BD , etc. Then the triangle AOF is rational (that is, has rational sides), since one of its sides, AF , is rational, and it is similar to the rational triangle ABD .



2678 [February, 1918]. Problem proposed by C. F. GUMMER, Queen's University, Canada.

Find necessary and sufficient conditions that the roots of the equation $x^{n+1} + a_1x^n + a_2x^{n-1} + \dots + a_{n+1} = 0$ may be all real and separated by the roots of $x^n + b_1x^{n-1} + b_2x^{n-2} + \dots + b_n = 0$.

SOLUTION BY THE PROPOSER.

Consider the equations

$$(1) \quad f(x) \equiv x^{n+1} + a_1x^n + \dots = 0,$$

$$(2) \quad g(x) \equiv x^n + b_1x^{n-1} + \dots = 0,$$

$$(3) \quad R_1(x) \equiv c_0x^{n-p} + c_1x^{n-p-1} + \dots = 0,$$

$$(4) \quad R_2(x) \equiv d_0x^{n-p-q} + d_1x^{n-p-q-1} + \dots = 0,$$

where $R_1(x)$ is the remainder with sign changed on dividing $f(x)$ by $g(x)$, $R_2(x)$ has the same relation to $g(x)$ and $R_1(x)$, etc.